

Characterization of Air Pollution Exposures as Risk Factors for Tuberculosis Infection

J. Lucian Davis, MD, MAS^{1,2,3}

William Checkley, MD PhD^{4,5}

¹Department of Epidemiology of Microbial Diseases and ²Center for Methods in Implementation and Prevention Science Yale School of Public Health , New Haven, CT USA

³Pulmonary Critical Care and Sleep Medicine, Yale School of Medicine, New Haven, CT USA

⁴Division of Pulmonary and Critical Care, School of Medicine, Johns Hopkins University, Baltimore, MD USA

⁵Center for Global Non-Communicable Disease Research and Training, School of Medicine, Johns Hopkins University, Baltimore, MD USA

Correspondence: J. Lucian Davis, MD, MAS

Yale School of Public Health - Epidemiology of Microbial Diseases

New Haven, Connecticut, 06520-8034

United States

E: Lucian.Davis@yale.edu

Running Head: Air pollution and latent tuberculosis

Word Count: 718 (Limit 1000)

References: 12 (Limit 15)

Tuberculosis is a leading cause of death due to infectious disease worldwide, accounting for 1.4 million deaths of the 10 million persons with active tuberculosis disease in 2019.¹ While improvements in the diagnosis and management of tuberculosis have led to declines in mortality over the past two decades, annual incidence remains unchanged. With the rapid growth in the world's population which currently includes approximately 1.7 billion persons with prevalent latent tuberculosis infection², a higher absolute number of incident tuberculosis cases progressing from latent infection threatens to offset gains from improved case detection and treatment success. At the first-ever United Nations High Level Meeting on Tuberculosis in 2018, new commitments were made to scaling up prevention measures. However, low rates of uptake and completion of tuberculosis preventive therapy and lack of highly effective vaccines highlight the need for new public health approaches to tuberculosis prevention.¹

One area of tuberculosis epidemiology that offers opportunities for improving prevention is enhancing our understanding of how exposure to environmental air pollutants may influence the risks of transmission and progression from latent infection to active disease.³ For example, occupational exposure to dust and smoke, including silica, are well-recognized risk factors for active tuberculosis disease.^{4,5} Second-hand exposure to tobacco smoke is also associated with a moderately higher risk of latent tuberculosis infection and a dose-dependent risk of active tuberculosis disease, especially among children.⁶ Exposure to solid fuel smoke and kerosene for heating and cooking are other potential factors that may increase the risk of active tuberculosis disease; however, results from epidemiological studies remain mixed⁷⁻¹⁰ and better studies are needed to understand if these are important risk factors. Even less is known about other air pollutants such as fine particulate matter (PM_{2.5}). At least one analysis of data from Beijing and Hong Kong has shown an association between seasonal concentrations of PM_{2.5} and active tuberculosis case notifications.¹¹

Here, Blount and colleagues present the results of a cross-sectional study examining associations between multiple urban environmental exposures and latent tuberculosis infection among 109 child household contacts of 72 patients with active tuberculosis in Hanoi, Vietnam.¹² Key indoor air pollution exposure variables were assessed by questionnaire and were complemented by measurement of personal exposures to PM_{2.5} over 48-72 hours. The authors constructed a multiple variable logistic regression model to explore the associations between these exposures and the outcome of latent tuberculosis infection, as defined by a positive tuberculin skin test.

This study makes some important observations. First, it confirms findings of a positive association between secondhand tobacco smoke exposure and latent tuberculosis infection. Second, the authors found that living on the ground floor was associated with a higher risk of latent tuberculosis infection when compared to floors above ground level. The authors conclude that indoor air pollution increases the risk of latent tuberculosis infection and suggest that mitigating these exposures may lead to better tuberculosis control.

There are some limitations to this study. While the assessment of personal exposures to PM_{2.5} was an important addition to this study, the lack of association with either questionnaire-based measurement of environmental risk factors or latent tuberculosis infection points to the complexity of air pollution assessment in epidemiological studies. First, air pollution assessment of urban exposures may require measurement of more than one pollutant (PM_{2.5}, black and elemental carbons, nitrogen oxides, carbon monoxide, sulfur dioxide, and ozone) to understand how air pollution exposures are associated with tuberculosis infection. Second, direct assessment of personal exposures over longer periods of time, combined with historical reconstruction of air pollution exposures over 2-5 years, may be required to improve our understanding of the relationship between air pollution and tuberculosis risk.

If air pollution increases transmission of tuberculosis, then what further actions are in order? First, there is a need for additional research to further characterize the sources, exposure intensities, and attributable tuberculosis risks associated with air pollution. Second, the causal mechanisms between environment and the host response to tuberculosis are not well-defined, and the findings in this study likely also generalize to other lung infections, including non-mycobacterial tuberculosis. Finally, these findings are a reminder to non-pulmonologists that tuberculosis is predominantly a pulmonary disease. Accordingly, assessing environmental exposures is an important part of both clinical history-taking and disease control, and should inform both individual counseling and population messaging about tuberculosis risk and prevention.

REFERENCES

1. World Health Organization. Global tuberculosis control: WHO Report 2020. Geneva: World Health Organization; 2020.
2. Houben RMGJ, Dodd PJ. The Global Burden of Latent Tuberculosis Infection: A Re-estimation Using Mathematical Modelling. *PLOS Medicine* 2016; **13**(10): e1002152.
3. Lienhardt C. From exposure to disease: the role of environmental factors in susceptibility to and development of tuberculosis. *Epidemiol Rev* 2001; **23**(2): 288-301.
4. Hnizdo E, Murray J. Risk of pulmonary tuberculosis relative to silicosis and exposure to silica dust in South African gold miners [published erratum appears in *Occup Environ Med* 1999 Mar;56(3):215-6]. *Occupational and Environmental Medicine* 1998; **55**(7): 496-502.
5. Rees D, Murray J. Silica, silicosis and tuberculosis. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease* 2007; **11**(5): 474-84.
6. Patra J, Bhatia M, Suraweera W, et al. Exposure to Second-Hand Smoke and the Risk of Tuberculosis in Children and Adults: A Systematic Review and Meta-Analysis of 18 Observational Studies. *PLOS Medicine* 2015; **12**(6): e1001835.
7. Pokhrel AK, Bates MN, Verma SC, Joshi HS, Sreeramareddy CT, Smith KR. Tuberculosis and indoor biomass and kerosene use in Nepal: a case-control study. *Environ Health Perspect* 2010; **118**(4): 558-64.
8. Sumpter C, Chandramohan D. Systematic review and meta-analysis of the associations between indoor air pollution and tuberculosis. *Tropical Medicine & International Health* 2013; **18**(1): 101-8.

9. Lin HH, Suk CW, Lo HL, Huang RY, Enarson DA, Chiang CY. Indoor air pollution from solid fuel and tuberculosis: a systematic review and meta-analysis. *The International Journal of Tuberculosis and Lung Disease* 2014; **18**(5): 613-21.
10. Emuron D, Siddharthan T, Morgan B, et al. Lack of an Association Between Household Air Pollution Exposure and Previous Pulmonary Tuberculosis. *Lung* 2019; **197**(6): 793-801.
11. You S, Tong YW, Neoh KG, Dai Y, Wang C-H. On the association between outdoor PM2.5 concentration and the seasonality of tuberculosis for Beijing and Hong Kong. *Environmental Pollution* 2016; **218**: 1170-9.
12. Blount RJ, Phan H, Trinh T, Dang H, Merrifield C, Zavala M, et al. Indoor Air Pollution and Susceptibility to Tuberculosis Infection in Urban Vietnamese Children. *Am J Respir Crit Care Med* [online ahead of print] 03 August 2021; <https://www.atsjournals.org/doi/abs/10.1164/rccm.202101-0136OC>.